arXiv:1207.0012v1 [quant-ph] 29 Jun 2012

# Semiclassical Coherent states path integrals

Alejandro M.F Rivas[*]
*Departamento de Física,*
*Comisión Nacional de Energía Atómica. Av. del Libertador 8250,*
*1429 Buenos Aires, Argentina.*

(Dated: July 3, 2012)

We have obtained a semiclassical approximation of the matrix elements of a quantum propagator for coherent sates basis, for that propose we have used the symplectically invariant Weyl representation of the propagator and gone beyond the common stationary phase approximation. The comparison with a "realistic" chaotic system that derives from a quadratic Hamiltonian, the cat map, reveals that the expression here derived is exact up to quadratic Hamiltonian systems.

Path integrals appear as a useful calculation tool for many quantum and statistical mechanical problems [1], while coherent sates are widely known to represent quantum states with the most classical resemblance, in the case of an harmonic oscillator they obey the classical equations of motion and are minimal uncertain states. Also, the coherent sates form an overcomplete basis which is a necessary ingredient for the construction of a path integral [2]. It implies the existence of several forms of path integrals, all quantum mechanically equivalent, but each leading to a different semiclassical limit. The formulation of path integrals applied to coherent sates has become widely used in many areas despite the very shaky mathematical background [3]. As was recently pointed out, the coherent states path integral breaks down in certain cases [4], where neither the action suggested by Weyl ordering nor the one constructed by normal ordering gives correct results when the Hamiltonian involves terms that are non linear in generators. In order to understand the quantum classical limit it is imperious to have a correct semiclassical expression of the quantum propagator in the most classical states, that is, in coherent states. In this Letter, by using a symplectically invariant formulation, we derive a semiclassical expression for the coherent states propagator that goes beyond the common stationary phase approximation [5]. The comparison with a "realistic" chaotic system that derives from a quadratic Hamiltonian for which the semiclassical limit is exact, the cat map, reveals that the formulation is exact up to quadratic Hamiltonian systems.

The coherent states $|X\rangle$ centered on points $X = (P, Q)$ in phase space, are obtained by translating to $X$ the ground state of the harmonic oscillator, its position representation is

$$\langle q\,|X\rangle = \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\frac{\omega}{2\hbar}(q-Q)^2 + i\frac{P}{\hbar}\left(q-\frac{Q}{2}\right)\right]. \tag{1}$$

For simplicity, unit frequency ($\omega = 1$) and mass ($m = 1$) are chosen for the harmonic oscillator without loss of generality. The overlap of two coherent states is then

$$\langle X\,|X'\rangle = \exp\left[-\frac{(X-X')^2}{4\hbar} - \frac{i}{2\hbar}X\wedge X'\right]. \tag{2}$$

Where the wedge product

$$X\wedge X' = PQ' - QP' = (\mathcal{J}X)\,.X',$$

In what follows we use the centers and chords formulation developed by Ozorio de Almeida [6] for both classical and quantum mechanics. The matrix elements of the unitary propagator $\widehat{U}^t$, that governs the quantum evolution of an $L$ degrees of freedom system, in the coherent states basis, $< X_1|\widehat{U}^t|X_2 > \equiv U^t(X_1, X_2)$ are

$$U^t(X_1,X_2) = \left(\frac{1}{\pi\hbar}\right)^L \int < X_1|U^t(x)\widehat{R}_x|X_2 > dx. \tag{3}$$

We have inserted the, symplectically invariant, Weyl decomposition of the propagator in terms of reflection operators [6]

$$\widehat{U}^t = \frac{1}{(\pi\hbar)^L}\int dx U^t(x)\widehat{R}_x, \quad \text{with} \quad U^t(x) = tr\left[\widehat{R}_x\widehat{U}^t\right], \tag{4}$$

with $\widehat{R}_x$ the set of reflection operators thought points $x = (p, q)$ in phase space [6, 9] then the operator $\widehat{U}^t$is written in terms of the function in phase space, $U^t(x)$, its center or Weyl Wigner symbol. The action of the reflection operator $\widehat{R}_x$ on a coherent state $|X\rangle$ is the $x$ reflected coherent state [8]:

$$\widehat{R}_x\,|X\rangle = e^{\frac{i}{\hbar}X\wedge x}\,|2x - X\rangle\,. \tag{5}$$

So that, inserting (5)and (2) in (3) the propagator in coherent states becomes,

$$U^t(X_1,X_2) = \frac{1}{(\pi\hbar)^L} e^{\frac{iX_1\wedge X_2}{2\hbar}} \int U^t(x) e^{\left[\frac{i}{\hbar}x\wedge\xi_0 - \frac{(\bar{X}-x)^2}{\hbar}\right]} dx. \tag{6}$$

with $\xi_0 \equiv (X_1 - X_2)$ the chord joining the coherent states points, and $\bar{X} = \frac{X_1+X_2}{2}$ is the mid point.

---

[*] member of the CONICET

For an Hamiltonian system, the propagator is $\widehat{U}^t = \exp[-i\frac{t}{\hbar}\widehat{H}]$ with $\widehat{H}$ the quantum Hamiltonian operator of the system. As has been seen in [6], that for sufficiently small times the Weyl symbol of such a propagator can be written as

$$U^t(x) = |\det[1 - (\mathfrak{J}\frac{t}{2}\mathcal{H})^2]|^{\frac{1}{4}} \exp\left[-\frac{it}{\hbar}H(x)\right] + O(t^2\hbar^2). \tag{7}$$

with $H(x) = tr\left[\widehat{R}_x\widehat{H}\right]$ the center or Weyl symbol of the Hamiltonian operator $\widehat{H}$ and where $\mathcal{H}$ is the corresponding Hessian matrix evaluated at the point $x$. In analogy to the classical theory, $H(x)$ will be within $O(\hbar)$ the classical Hamiltonian $H_c(x)$. Also, if $H(x)$ is quadratic (7) represents a unitary operator .

For longer times $t$, the composing an even number $K$ of unitary operators for times $\varepsilon = t/K$ is performed. In the Weyl representation such composition results [6]

$$U^t(x) = \int \frac{dx_i^K}{(\pi\hbar)^{KL}} \prod_{j=1}^{K} U^{\frac{t}{K}}(x_j) e^{\frac{i}{\hbar}\Delta_{K+1}(x,x_1,\cdots,x_K)}, \tag{8}$$

with $dx_i^K = dx_K \cdots dx_1$ and $\Delta_{K+1}(x, x_1, \cdots, x_K)$ the symplectic area of the polygon with endpoints centered on $x$ and whose $j'$th side is centered on $x_j$. This formula is exact, while in the limit $K \to \infty$, the expression (7) can be inserted for $U_{\frac{t}{K}}(x)$. In which limit the amplitude vanishes, yielding to the *path integral*

$$U^t(x) = \lim_{K\to\infty} \int \frac{dx_i^K}{(\pi\hbar)^{KL}} e^{\frac{i}{\hbar}\left[\Delta_{K+1}(x,x_1,\cdots,x_K) - \frac{t}{K}\sum_{j=1}^{K} H(x_j)\right]}. \tag{9}$$

Analogously, for the case of classical composing of an even number $K$ flows for time $\varepsilon = t/K$ in the limit where $K \to \infty$, the center generating function of the canonical transformation is [6]

$$S^t(x) = \lim_{K\to\infty} -\frac{t}{K}\sum_{j=1}^{K} H_c(x_j) + \Delta_{K+1}(x, x_1, \cdots, x_K). \tag{10}$$

The polygon $\Delta_{K+1}$ has one large side $\xi$ passing through the center $x$ and $K$ small chords tangents to the orbit as $K \to \infty$ such that $\partial S^t/\partial x_j = 0$. Hence, the *center variational principle* is obtained: The *center action*

$$S_\gamma^t(x) = \oint_x p \cdot dq - \int_\gamma H_c(x)dt \tag{11}$$

is stationary along the classical trajectory $\gamma$. The paths to be compared always have their endpoints centered on the point $x$. The second integral is evaluated along this path, whereas the first integral is closed off by the chord centered on $x$.

Hence, the phase of the integral in (9) coincides with the center action $S_\gamma^t(x)$ for the polygonal path $\gamma$. The center variational principle ensures that this center action is stationary for the classical trajectories centered on $x$. The semi-classical approximation for the Weyl propagator was obtained by performing a stationary phase approximation in (9) then the *center variational principle* yields [6] :

$$U^t(x)_{SC} = \sum_\gamma \frac{2^L \exp\left\{i\hbar^{-1}S_\gamma^t(x) + i\alpha_\gamma^t\right\}}{|\det(\mathcal{M}_\gamma + 1)|^{\frac{1}{2}}}, \tag{12}$$

the summation is over all the classical orbits $\gamma$ whose center lies on the point $x$ after having evolved a time $t$ [6]. Where $S_\gamma^t(x)$ is the classical center generating function of the orbit, $\mathcal{M}_\gamma$ is the symplectic matrix for the linearized transformation between the neighborhood of the tips of the chord $\xi_\gamma$ joining the initial and final point of the orbit. For sufficiently short times such that the variational problem has an unique solution there will have a single chord. Although for longer times, there will be bifurcations producing more chords. In the case of a single orbit, the corresponding *Morse index* $\alpha_\gamma^t = 0$.

In order to obtain a coherent state path integral, expression (9) is inserted in the coherent state propagator (6) so that

$$U^t(X_1, X_2) = \lim_{K\to\infty} \int \frac{dx_i^K}{(\pi\hbar)^{KL}} e^{\frac{i}{\hbar}\left\{\frac{X_1\wedge X_2}{2} - \frac{t}{K}\sum_{n=1}^{K} H(x_n)\right\}} \times \int \frac{dx}{(\hbar\pi)^L} e^{\left\{\frac{i}{\hbar}[x\wedge\xi_0 + \Delta_{K+1}(x,x_1,\cdots,x_K)] - \frac{(\bar{X}-x)^2}{\hbar}\right\}} \tag{13}$$

Recalling the linear relation of $\Delta_{K+1}$ with $x$ ,

$$\Delta_{K+1}(x, x_1, \cdots, x_K) = C + \xi \wedge x, \text{ with } \xi = \sum_{j=1}^{K}(-1)^j x_j. \tag{14}$$

The $x$ integral in (13) is quadratic and can then be solved exactly yielding

$$U^t(X_1, X_2) = e^{-\frac{iX_1\wedge X_2}{2\hbar}} \lim_{K\to\infty} \int \frac{dx_i^K}{(\pi\hbar)^{KL}} e^{-\frac{1}{4\hbar}(\bar{\xi}-\xi_0)^2} \times e^{\frac{i}{\hbar}\left\{\Delta_{K+1}(\bar{X},x_1,\cdots x_K) - \frac{t}{K}\sum_{n=1}^{K} H(x_n)\right\}} \tag{15}$$

where $\bar{\xi}$ is the chord passing through the mid point $\bar{X}$. Note that according to (14) the chord $\bar{\xi}$ depends only on the other centers $x_1, \cdots, x_K$ . The usual semiclassical limit of the coherent states propagator is obtain, in the limit where $\hbar \to 0$ , for a stationary phase approximation of (15). The phase in (15) is the center generating function (10). Hence, according to the center variational principle the path integral is stationary for classical orbits, then analogously to (12) stationary phase approximation yields

$$U^t(X_1, X_2)_{SC1} = \sum_\gamma \frac{2^L e^{\left\{\frac{i}{\hbar}\left[\widetilde{S}_\gamma^t(\bar{X}) - \frac{1}{2}X_1\wedge X_2\right] - \frac{1}{4\hbar}\left(\bar{\xi}_\gamma - \xi_0\right)^2\right\}}}{|\det(\mathcal{M}_\gamma + 1)|^{\frac{1}{2}}} \tag{16}$$

where the index runs over all the contributing classical orbits $\gamma$ with center generating function$S^t_\gamma(\bar{X})$ whose center lies on the point $\bar{X}$ and $\bar{\xi}_\gamma$ is the chord joining the initial and final point of the orbit. Again, will have a single chord for sufficiently short times.However, as a consequence of a Gaussian cutoff on the length of the chords, the amplitude will be severely damped if the classical chord $\bar{\xi}_\gamma$ is long. We have define the action $\widetilde{S}^t_\gamma(x) = S^t_\gamma(x) + \hbar\alpha^t_\gamma$ in order to include the Morse index in the action. Equation (16) has been obtained by an equivalent procedure by dos Santos and de Aguiar [5]. Also, as will be shown in Fig 1. expression (16) does not give accurate results for quadratic Hamiltonian systems.

An other way to obtain a semiclassical approximation for the coherent sates propagator is to insert the semiclassical approximation for the Weyl propagator (12) in (6) so that

$$U^t_{SC}(X_1, X_2) = \left(\frac{2}{\pi\hbar}\right)^L e^{\frac{i}{2\hbar}X_1\wedge X_2} \sum_\gamma e^{i\alpha^t_\gamma} \\ \times \int \frac{\exp -\frac{1}{\hbar}\left(\overline{X} - x\right)^2}{|\det\left(\mathcal{M}_\gamma + 1\right)|^{\frac{1}{2}}} e^{\frac{i}{\hbar}\left[S^t_\gamma(x) + \xi_0\wedge x\right]} dx \tag{17}$$

This procedure is equivalent to perform in (13) the stationary phase approximation in the $dx_n \cdots dx_1$ variables before integrating in $dx$. We must now perform the $x$ integral, in a first place let us perform an ordinary stationary phase approximation in (17). Expanding the phase of (17) around the stationary point $x_0$ and performing an usual stationary phase approximation, another semiclassical approximation for the propagator in coherent states rises

$$U^t(X_1, X_2)_{SC2} = \sum_\gamma \frac{2^L e^{\left\{\frac{i}{\hbar}\left[\widetilde{S}^t_\gamma(\xi_0) + \frac{1}{2}X_1\wedge X_2\right] - \frac{1}{\hbar}\left(x_0 - \bar{X}\right)^2\right\}}}{|\det(\mathcal{M}_\gamma - 1)|^{\frac{1}{2}}} \tag{18}$$

where $x_0$ must satisfy $\xi(x_0) = -\mathcal{J}\frac{\partial S^t_\gamma(x)}{\partial x}\Big|_{x=x_0} = \xi_0$, that is the chord through $x_0$ equals the chord joining the coherent sates, and the chord action $\widetilde{S}^t_\gamma(\xi_0) = \widetilde{S}^t_\gamma(x_0) - \xi_0 \wedge x_0$ [6] being the Legendre transform of $\widetilde{S}^t_\gamma(x_0)$. This expression (18) is complementary to equation (16), it is written in terms of chords instead of centers. Although, as is the case for the chord action [6], the expression (18) diverges for very short times where the monodromy matrix becomes the identity.

In obtaining (18) it was assumed that the Gaussian term in (17) is smooth close to the stationary point. Although, there is a compromise between the point of stationary phase and the maximum of the Gaussian in $\bar{X}$ . If a stationary phase point falls far from $\bar{X}$ its contribution will be severely damped. For this reason, it is not surprising that expression (18) fails, as is shown in Fig 1. Also, it must be noted that classical orbits that starts near $X_2$ and ends near $X_1$ will have an important contribution, this orbits will have a center point close to $\bar{X}$. Hence, in order to perform the phase space integral in (17), let us expand the center action up to quadratic terms near the mid point $\bar{X}$ so that

$$S^t_\gamma(x) = S^t_\gamma(\bar{X}) + \bar{\xi}\wedge x' + x' B_{t\gamma} x' + O(x'^3) \tag{19}$$

with $x = \bar{X} + x'$ and $S^t_\gamma(\bar{X})$ is the action of the orbit through the mid point $\bar{X}$ , while the symmetric matrix $B_{t\gamma}$ is the Cayley representation of $\mathcal{M}_\gamma$

$$\mathcal{J}B_{t\gamma} = \frac{1 - \mathcal{M}_\gamma}{1 + \mathcal{M}_\gamma} = \frac{1}{2}\mathcal{J}\frac{\partial^2 S^t_\gamma(x)}{\partial x^2}. \tag{20}$$

After the linearization of the flux around the mid point (19) is used in (17) a quadratic integral is obtained that can be exactly solved. Using

$$I = \int e^{\frac{1}{\hbar}\left[-x' V_t x' + Y.x'\right]} dx' = \frac{(\pi\hbar)^L e^{\left[\frac{1}{4\hbar}Y V_t^{-1} Y\right]}}{\sqrt{(\det V_t)}}, \tag{21}$$

the semiclassical propagator in coherent states, takes the form

$$U^t(X_1, X_2)_{SC3} = \sum\nolimits_\gamma \frac{2^L}{|\det[V_t(\mathcal{M}_\gamma + 1)]|^{\frac{1}{2}}} \exp\left[-\tfrac{1}{\hbar}\delta\overline{C}_t\delta\right] \\ \times \exp\tfrac{i}{\hbar}\left[-\tfrac{1}{2}X_1\wedge X_2 + \widetilde{S}^t_\gamma(\bar{X}) + \delta\overline{B}_t\delta + \hbar\tfrac{\varepsilon}{2}\right], \tag{22}$$

with the complex matrix $V_t = 1 - iB_{t\gamma}$, such that $\det V_t = |\det V_t|e^{i\varepsilon}$, the point shift $\delta = {}^1\!/_2\left(\bar{\xi} - \xi_0\right)$, while the real matrices $\overline{C}_t$ and $\overline{B}_t$ are defined such that

$$\overline{C}_t - i\overline{B}_t = \left(1 + i\mathcal{J}B_{t\gamma}\mathcal{J}\right)^{-1}. \tag{23}$$

The expression (22) of the semiclassical matrix elements between two coherent states of the quantum propagator, is the main contribution of this Letter. It is expressed entirely in terms of classical objects, namely the action $S^t_\gamma(\bar{X})$ of the classical orbit whose mid point is $\bar{X}$ , the point shift $\delta$, the monodromy matrix $\mathcal{M}_\gamma$ and its Cayley representation $B_{t\gamma}$. It is important to remark the Gaussian term that dampens the amplitude for large values of the point shift $\delta$, that is for orbits that starts far from the points $X_2$ (or end far from $X_1$ ). So as, the main contribution in the sum over classical orbits in (22) will come from the orbits $\gamma$ , whose initial point $x_i$ lies the closest to $X_2$. It is important to remark that, the same expression (22) could have obtain by perform in (15) the complete quadratic integral instead of a stationary phase approximation that led to (16).

Also, note that, if $t = 0$, the quantum propagator is just the identity operator in Hilbert space, the classical symplectic matrix is the identity, the center action is null, and the result (2) for the overlap of coherent sates is recovered

We will study the matrix elements on a surface of section that is transversal to the flux, in analogy with classical Poincaré surfaces of section. The study of autonomous fluxes through a map on surface of section is a standard procedure, and quantum surface of section methods are shown to be exact [10] for general Hamiltonian systems. Here, the theory is applied to the cat map i.e. the linear automorphism on the 2-torus generated by the $2 \times 2$ symplectic matrix $\mathcal{M}$, that takes a point $x_-$ to a point $x_+ : x_+ = \mathcal{M}x_- \mod(1)$. With the symplectic matrix

$$\mathcal{M} = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}. \tag{24}$$

This map is known to be chaotic, (ergodic and mixing) as all its periodic orbits are hyperbolic, it corresponds to viewing stroboscopically the motion generated by a quadratic Hamiltonian [7]. Quantum mechanics on the torus, implies a finite Hilbert space of dimension $N = \frac{1}{2\pi\hbar}$, and that positions and momenta are defined to have discrete values in a lattice of separation $\frac{1}{N}$ [9, 11]. The cat map was originally quantized by Hannay and Berry [11] in the coordinate representation the propagator is:

$$< \mathbf{q}_k|\hat{\mathbf{U}}_{\mathcal{M}}|\mathbf{q}_j >= \left(\frac{i}{N}\right)^{\frac{1}{2}} e^{\frac{i2\pi}{N}(k^2-jk+j^2)}, \tag{25}$$

where the states $< q|\mathbf{q}_j >$ are periodic combs of Dirac delta distributions at positions $q = {}^{j}/_{N} mod(1)$, with $j$ integer in $[0, N-1]$. Bold fonts characters are used to describe quantum objects on the torus. The definition of the periodic coherent state [9], in accordance to (1) is

$$< \mathbf{X}|\mathbf{q}_k >= \sum_{j=-\infty}^{\infty} e^{-\frac{1}{\hbar}\left[iP(j+\frac{Q}{2}-{}^{k}/_{N})+\frac{1}{2}(j+Q-{}^{k}/_{N})^2\right]}. \tag{26}$$

Indeed the construction on the torus from the plane is obtain in terms of averages of translations [9].

Figure 1 shows the relative error $E$ of the amplitude of the semiclassical approximations $U^t(X_1, X_2)_{SC1}$ and $U^t(X_1, X_2)_{SC2}$ obtained in (16) and (18) with respect to the exact expressions obtained with the quantum propagator (25) on coherent states (26). It can be seen that neither (16) nor (18) are good approximations of the exact coherent sates matrix elements giving errors in the amplitude of more than 10% or that highly grows with $N$ respectively. While we have verified that $U^t(X\quad X\;)$



Figure 1: Relative error of the amplitude of the Semiclassical expressions $U^t(X_1, X_2)_{SC1}$ and $U^t(X_1, X_2)_{SC2}$ as a function of $N$. In dotted line error of $U^t(X_1, X_2)_{SC1}$ and in full line the error of $U^t(X_1, X_2)_{SC2}$.

To conclude, in this work the semiclassical coherent state propagator has been obtained using the Weyl representation. The expression of $U^t(X_1, X_2)_{SC3}$ obtained in (22) goes beyond a stationary phase approximation and is exact for linear systems. Also, the comparison with a system whose semiclassical limit is exact has allows to correctly check the exactness of the obtained expression up to quadratic Hamiltonian systems.

I thanks G. Carlo, E. Vergini and M. Saraceno for stimulating discussions and the CONICET for financial support.

---

[1] J.R. Klauder, in *Path Integrals and their Applications in Quantum, Statistical and Solid State Physics*, edited by G. Papadopoulos and J. Devreevese. (Plenum Press, Antwerp, Belgium, 1977)

[2] J.R. Klauder and B.S. Skagerstam, *in Coherent Sates: Aplications in Physics and Mathematical Physics*, (Singapore: Word Scientific, 1985).

[3] J.R. Klauder, arXiv:quant-ph/0303034.

[4] J. H. Wilsom and V. Galitski, Phys. Rev. Lett **106**, 110401 (2011).

[5] L.C. dos Santos and M.A.C. de Aguiar, J. Phys. A: Math. Gen. **39**, 13465 (2006).

[6] A.M. Ozorio de Almeida, Physics Report **295**, 266 (1998).

[7] J. P. Keating ,Nonlinearity **4**, 277 (1991).

[8] A.M.F. Rivas, J. Phys. A: Math. Gen. **40**, 11057 (2007).

[9] A.M.F. Rivas and A.M. Ozorio de Almeida, Annals of Physics **276,** 223 (1999).

[10] T. Prosen, J. Phys. A: Math. Gen. **28**, 4133 (1995).

[11] J.H. Hannay and M.V. Berry, Physica **D**, 267(1980).